



(12)

EUROPEAN PATENT SPECIFICATION

- (43) Date of publication of patent specification: **01.09.93** (51) Int. Cl.⁵: **A61K 31/56**, C07J 1/00, C07C 49/225, C07C 33/34, C07C 69/017, C07C 39/17, C07J 63/00, C07C 49/237, C07D 303/04, C07C 33/36, A61K 31/045
- (21) Application number: **89310063.6**
- (22) Date of filing: **02.10.89**

(94) Anti-androgen compounds.

- | | |
|--|---|
| <p>(30) Priority: 30.09.88 US 252375
27.09.89 US 413249</p> <p>(43) Date of publication of application:
11.04.90 Bulletin 90/15</p> <p>(43) Publication of the grant of the patent:
01.09.93 Bulletin 93/35</p> <p>(94) Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE</p> <p>(56) References cited:
FR-A- 1 326 731
FR-M- 6 828
GB-A- 1 119 068</p> <p>JOURNAL OF MEDICINAL CHEMISTRY, vol. 15, no. 12, December 1972, pages 1336-1338, Washington, DC; US; A. P. ROSZKOWSKI et al, "Effects of some substituted phenanthrenes on the central nervous system"</p> | <p>(73) Proprietor: ARCH DEVELOPMENT CORPORATION
1101 East 58th Street
Chicago, Illinois 60637(US)</p> <p>(72) Inventor: Liao, Shutsung
5632 South Woodlawn Avenue
Chicago, Illinois 60637(US)
Inventor: Harvey, Ronald, G
5841 South Maryland Avenue Box 424
Chicago, Illinois 60637(US)
Inventor: Pataki, John
5440 Kimbark Avenue
Chicago, Illinois 60615(US)</p> <p>(74) Representative: Brown, John David et al
FORRESTER & BOEHMERT Franz-
Joseph-Strasse 38
D-80801 München (DE)</p> |
|--|---|

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid (Art. 99(1) European patent convention).

JOURNAL OF ORGANIC CHEMISTRY, vol. 53,
no. 18, 2nd September 1988, pages
4253-4256, Washington, DC, US; H. LEE et al,
"New synthetic approaches to cyclopenta(a)
phenanthrenes and their carcinogenic de-
rivatives"

Description

The present invention relates generally to new classes of anti-androgen compounds, their method of synthesis and their use as anti-androgens.

5 Androgens are one of the five major classes of steroid hormones. Steroid hormones form complexes with receptor proteins which are distributed in a tissue specific fashion within target cells. Jensen, et al., Proc. Nat'l. Acad. Sci. (USA), 59:632 (1968); Gorski, et al., Ann. Rev. Physiol., 42:17 (1976); and Liao, et al., page 633 in Biochemistry of Hormones, H.L.J. Makin, ed. (Blackwell Sci. Publ. Oxford, 1984). Investigation of the specificity and affinity of steroid hormones for their particular cellular receptors(s) has contributed
10 greatly to the understanding of the relationship between structure and biological activity, target organ specificity, overlaps in hormonal activity, and the mechanism of action of many anti-hormones. Liao, S., et al., J. Biol. Chem., 248:6154 (1973); Liao, S., Biochemical Actions of Hormones (Litwack, G., ed), 4:35T, Academic Press, New York (1977); Liao, et al., Biochemistry of Steroid Hormones, (Makin, H.L.J., ed.) pp. 630-680, Blackwell Scientific Publications, Oxford (1984).

15 Studies on the structural recognition of ligands by steroid receptors suggests that compounds with geometric structures similar to that of natural androgens, such as 5 α -dihydrotestosterone, can bind tightly to androgen receptors and can act either as potent androgens or as anti-androgens. Liao, S., et al., J. Biol. Chem., 248:6154 (1973). The most well known steroidal anti-androgens are cyproterone and its acetate [- (Neumann, Androgens and Anti-androgens, (L. Martini and M. Motta, ed.) pp. 163, Raven Press, New York
20 (1977)] which act by interacting with androgen receptors and prevent androgens from binding to the receptors. Fang, S., Molec. Pharmacol., 5:428 (1969). Non-steroidal anti-androgens, such as flutamide-related compounds, also act through the same mechanism. Liao, S., et al., Endocrinol., 94:1205 (1974) and Neri, R., Androgens and Anti-androgens, (L. Martini & M. Motta, ed.) pp. 179, Raven Press, New York (1977).

25 Studies of the topographic recognition of cyclic hydrocarbons and related compounds by receptors for various steroid hormones have suggested that while the hormonal action of a steroid may be dependent upon the interaction of a functional group present on the hormone with a specific group present on the receptor, the presence of such a functional group may not be required for the antagonistic activities of a compound that can physically block hormone binding to the receptor. Thus, many small molecules, that were hitherto considered to be biologically inert, may interact with steroid receptors specifically and affect
30 hormonal activities in vivo. Chang, C., et al., J. Steroid Biochem., 27:123 (1987). One such example is 9,10-dihydrophenanthrene (DHP) which, in comparison to a natural androgen, lacks one of the rings as well as two functional groups, including a double-bonded oxygen and a hydroxyl group. Despite these differences, DHP nevertheless interacts with androgen receptors in cell-free systems and inhibits the growth of the ventral prostate in rats. Chang, C., et al., J. Steroid Biochem., 27:123 (1987).


35 Lee, H. and Harvey R.G., J. Org. Chem. 1988 53, pp.4253-4256 discloses 8,7,16,17-tetrahydro-15H-cyclopenta[a]phenantren-17-one, as well as 16,17-dihydro-15H-cyclopenta[a]phenantren-17-one.

Roszkowski, A.P. et Al. J. Med. Chem. 1972, 15, pp.1336-1338 discloses 2-(2-hydroxyisopropyl)-phenantrene as well as 2-(2-hydroxyisopropyl)-9,10-dihydrophenantrene and their use as paralytics, analgesics and anticonvulsants. not used in the United States of America because of their side effects, including
40 undesired hormonal activities and/or toxicities.

Thus, there continues to exist a need in the art for new classes of anti-androgens which do not have other hormonal activities and/or side effects, yet, which are useful in treating diseases or abnormalities related to androgen responsive organs.

BRIEF SUMMARY OF THE INVENTION

The present invention relates generally to an anti-androgen comprising a compound selected from the group consisting of compounds of the general formula:



wherein said ring A is substituted or unsubstituted and wherein said ring A is substituted it is fused at either the 1-2 or 2-3 positions to a hydroxycycloalkyl, or a ketocycloalkyl, or is substituted with a hydroxyl, an alkyl, or a hydroxyalkyl other than hydroxyisopropyl other than hydroxyisopropyl at the 1, 2 or 3 positions;

wherein said ring B is saturated or mono-unsaturated at the 9-10 position and is optionally substituted with a lower alkyl at either the 9 or 10 position;

wherein said ring C is substituted or unsubstituted, and wherein when said ring C is substituted it is fused at the 6-7 or 7-8 positions to a ketocycloalkyl or a hydroxycycloalkyl; and

provided that rings A and C may not both be unsubstituted, wherein the compound is not 6,7,16,17-tetrahydro-15H-cyclopenta(a)phenanthren-17-one or 16, 17-dihydro-15H-cyclopenta(a)phenanthren-17-one, or a derivative of said compound wherein a free hydroxyl group has been replaced with an acetate or a propionate moiety.

Preferably, the ketocycloalkyl is a cyclopentanone or cyclohexanone; the hydroxycycloalkyl is cyclopentanol or cyclohexanol, and the alkyl is methyl.

Preferred derivatives include 1-[2-(9,10-dihydrophenanthryl)]-1-ethanol; 6,7,16,17-tetrahydro-15H-cyclopenta[a]-phenanthren-17-ol; 1-[2-(9(or 10)-methyl-9,10-dihydrophenanthryl)]-1-ethanol; 2-hydroxy-9,10-dihydrophenanthrene; 4'-oxo-7,8-cyclohexeno-9,10-dihydrophenanthren-1-ol; and 4'-oxo-7,8-cyclohexenopenanthren-1-ol. Also provided by the invention are the corresponding derivatives wherein a free hydroxyl group has been replaced with an acetate or a propionate moiety.

Also provided by the invention are pharmaceutical compositions comprising pharmaceutically effective amounts of one or more compounds of the invention in combination with a pharmaceutically acceptable solvent, diluent, adjuvant, or carrier,

In one preferred embodiment of the present invention, a therapeutically effective amount of one or more of the compounds of the invention can be administered to treat various disorders including those conditions wherein excessive androgenic activities have been implicated in the pathogenesis of certain androgen dependent conditions, such as benign prostatic hyperplasia, prostate cancer, male pattern baldness, female hirsutism, and the like. These pharmaceutical compositions, comprising compounds of the invention, can be administered by topically or internal routes.

In addition, it is expected that some of the compounds of the invention may interact with mutated receptors and, therefore, may be useful in the treatment, as well in the diagnosis of, androgen and other hormone-insensitive tumors. It is also expected that compounds of the invention will be important in the studies of the mechanism of action of androgens and anti-androgens. Thus, also provided by the invention are methods for localizing androgen receptors in, for example, tissue samples, wherein the sample is incubated with a compound of the invention and wherein the compound is labelled with a marker, the marker is detected and the androgen receptor in the sample is thereby localized.

Other aspects and advantages of the present invention will be apparent upon consideration of the following detailed description thereof which includes numerous illustrative examples of the practice of the invention, reference being made to the drawing wherein:

Figure 1 is a schematic representation of the synthesis of compounds 1, 3, 4, 5, 6, and 7;

Figure 2 is a schematic representation of the synthesis of compounds 8, 9, 10, 11, 12, and 13; and

Figure 3 is a schematic representation of the synthesis of compounds 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, and 24.

Figure 4 is a schematic representation of the synthesis of compounds 26, 27, 28, and 29.

DETAILED DESCRIPTION

The following examples illustrate practice of the invention in the synthesis of anti-androgen compounds and their activity and use as anti-androgens. More specifically, Example 1 relates to the synthesis of the compounds of the invention and Example 2 relates to a comparison of the relative binding activities of the compounds of the invention.

The examples which follow are for illustrative purposes only and are not intended in any way to limit the scope of the invention.

EXAMPLE 1Structural Considerations and Synthesis of Dihydrophenanthrene Derivatives

Among the requirements considered in constructing the new classes of anti-androgens were the following. Preferably, the geometry of the ligand molecule should be the same as, or smaller, than natural androgens, such as 5 α -dihydrotestosterone, so that the molecule can be easily inserted into the ligand binding site of the receptor and can have a high rate of association. In contrast to natural steroid hormones that have a rigid structure, the molecule preferably should have a flexible structure allowing it to conform to the structure of the receptor binding cavity and to become trapped. This results in a ligand with a high binding affinity and with a slow rate of dissociation. It is also preferable, but possibly not essential, that the molecule have a functional group present, at one or both ends of the molecule, that corresponds to the carbonyl group of ring-A and/or the hydroxyl group of ring-D of a natural androgen. It is also preferable that the molecule be able to interact or to bind to the receptor in a cell-free assay, in an organ or cell-binding assay, or in an *in vivo* biological assay.

A number of classes of compounds were synthesized following the above guidelines. These general classes include dihydrophenanthrene derivatives and chrysene derivatives. Their synthesis is described below. See also, Lee, H., *et al.*, *J. Org. Chem.*, 53:4253 (1988) and Harvey, R., *et al.*, *J. Org. Chem.*, 51:1407 (1986).

Shown in Figure 1 is a schematic representation of the synthesis of intermediates 1, 3, 4 and dihydrophenanthrene derivatives 5, 6 and 7, as described below.

2-(1-Naphthyl)ethyl iodide (1)

To a solution of P₂I₄ (7.778 g; 13.82 mmol) in CS₂ (350 ml) was added at once 2-naphthalen-ethanol. The mixture was stirred for 48 h. The solvent was evaporated and the residue was dissolved in ether. The ether solution was extracted with water 3 times. Column chromatography on Florisil™ afforded (1) as an oil (10 g, 67%).

2-[2-(1-Naphthyl)ethyl]cyclopentanone (3)

To a solution of (2) N-cyclohexyl-N-cyclopentenyl magnesium bromide, prepared from the reaction of N-cyclopentylidenecyclohexylimine [See Whitesell, J.K., Whitesell, M.A., *Synthesis*, page 517 (1983); and Stork, G., *et al.*, *J. Am. Chem. Soc.*, 85:2178 (1963)] (8.5 g, 415 mmol) with ethylmagnesium bromide (60 mmol; 10 mL of 3 mole solution in THF) was added 2-(1-naphthyl)ethyl iodide (1) (12 g; 42 mmol), and the resulting mixture was refluxed for 20 h. Hydrolysis was effected by refluxing with 180 mL of 10% aqueous HCl for 3 h. The product was extracted and further purified by chromatography on a column of Florisil™, Floridin Co., Florida. Elution with benzene afforded (3) (9.68 g; 96%) as an oil: NMR, δ 1.3-2.5 (m, 8, aliphatic), 3.1 (apparent, 2, benzylic), 7.2-8.2 (m, 7, aromatic). Anal. Calc'd. for C₁₇H₁₈O: C, 85.67; H, 7.61; Found: C, 85.72; H, 7.63.

16,17-Dihydro-15H-cyclopenta[a]phenanthrene (4)

Cyclization of (3) (6.4 g; 26 mmol) was carried out in polyphosphoric acid (10 ml) at 110°C for 2 h under N₂. Ice water was added, and the mixture was extracted with CH₂Cl₂. The product was worked up conventionally and chromatographed on a column of Florisil™ to yield a mixture of (4) and other products of acidic disproportionation. This mixture was dehydrogenated by heating with 10% Pd/C (1.8 g) in triglyme (250 ml) at reflux for 2 h under N₂. The reaction mixture was cooled and filtered, and the filtrate was diluted with ether and washed with water several times to remove triglyme. The ether solution was

dried over MgSO_4 and evaporated to dryness to afford a white solid which was triturated with cold hexane to yield pure (4) (14.9 g, 85%); mp 134-135 °C; NMR 2.27 (m, 1, $\text{H}_{2,6,7}$), 7.72 (s, 2, $\text{H}_{6,7}$), 7.85 (m, 1, H_8 , 8.50 (d, 1, $\text{H}_{1,12}$ = 8.2 Hz), 8.64 (m, 1, H_1).

6,7,16,17-Tetrahydro-15H-cyclopenta[a]phenanthrene (5)

Compound (4) (4.3 g; 19 mmol), dissolved in ethyl acetate (100 ml), was hydrogenated over a 10% Pd/C catalyst (3.2 g) at 50 lb/sg.i. pressure at room temperature for 24 h. Crystallization from hexane gave compound (5) as a white solid (2.97 g, 71%); m.p. 64-65°C; NMR; δ 2.2 (m, 2, aliphatic), 2.7-3.2 (m, 8, benzylic), 7.1-7.4 (m, 4, aromatic), 7.6 (d, 1, $\text{H}_{1,11}$), 7.7 (m, 1, H_1).

6,7,16,17-Tetrahydro-15H-cyclopenta[a]phenanthren-17-one (6)

The hydrocarbon (5) 48 mg; 0.22 mmol) was dissolved in hot acetic acid (15 ml) and water (5 ml) was added slowly with stirring to maintain homogeneity of the solution. 2,3-Dichloro-5,6-dicyano-1,4-benzoquinone (DDQ) (198 mg, 0.88 mmol) was added, changing the color of the solution to dark green. Stirring was continued at reflux for 30 min. during which time the color of the solution changed to dark red. The reaction mixture was cooled and diluted with ether, and the ether layer was washed with water and aqueous NaOH. The solution was dried over MgSO_4 and evaporated to provide a white solid which was chromatographed on a column of Florisil™. Elution with ether yielded 6 (39 mg, 76%); m.p. 127-128 °C; NMR δ 2.72 (t, 1, H_{15} or ϵ), 2.85-2.91 (m, 4, $\text{H}_{6,7}$), 3.07 (t, 1, H_{15} or ϵ), 7.24-7.31 (m, 3, $\text{H}_{2,3,4}$), 7.67-7.77 (m, 3, $\text{H}_{1,11,12}$). Anal. Calcd. for $\text{C}_{17}\text{H}_{14}\text{O}$: C, 87.25; H, 6.02. Found: C, 87.13; H, 6.03.

6,7,16,17-Tetrahydro-15H cyclopenta[a] phenanthren-17-ol (7)

A suspension of ketone (6) (300 mg) and NaBH_4 (200 mg) in MeOH (500 ml) and THF (30 ml) was stirred at room temperature for 2.5 h. Solvents were evaporated and the residue was taken up in ether. The ether solution was washed with H_2O three times. Column chromatography on Florisil™ afforded (7) (290 mg; 298%) as a solid, NMR; δ 2.8 (m, 8, $\text{H}_{6,7,15,15}$), 7.8 (m, 6, aromatics).

Shown in Figure 2 is a schematic representation of the synthesis of dihydrophenanthrene derivatives 8, 9, 10, 11, 12, and 13, as described below.

2-Acetyl-9,10-dihydrophenanthrene (8)

To a solution of 18 g (0.1 mole) 9,10-dihydrophenanthrene in 200 mL of dry CH_2Cl_2 , cooled in ice, 20 g (0.15 mole) of anh. AlCl_3 was added. To this solution 9.42 g (0.12 mmole) of acetyl chloride in 50 mL of dry CH_2Cl_2 was added in 17 mins. Stirring was continued for 30 mins in ice and 2 h at room temperature. The solution was poured on ice and 25 mL of concentrated HCl and stirred for 15 mins. The yellow organic layer was separated, washed with NaHCO_3 solution, and dried. The solvent was removed and the residue distilled at 0.15 mm pressure. The distillate, 15.28 g, crystallized from ethanol-hexane, m.p. 58-60 °.

1-[2-(9,10-Dihydrophenanthryl)]-1-ethanol (9)

To a stirred solution of 1.0 g of 2-acetyl-9,10-dihydrophenanthrene in 35 mL of MeOH was added 1 g of NaBH_4 in small portions within 45 mins. The solution was stirred for 2 h more. Most of the MeOH was removed under reduced pressure. Water was added to the residue, and the residue was extracted twice with CH_2Cl_2 ; the combined two extracts were dried and evaporated to dryness. The residue crystallized from ether, m.p. 84-85.5 °.

2-Acetyl-9(or 10)-methyl-9,10-dihydrophenanthrene (10)

To a stirred solution of 5.216 g (26.86 mmol) of 9-methyl-9,10-dihydrophenanthrene in 100 mL of dry CH_2Cl_2 , cooled in an ice-salt bath, 6.67 g (50 mmol) of anh. AlCl_3 was added. To the deep red solution, a solution of 2.2 mL (2.402 g; 32.23 mmol) of acetyl chloride in 25 mL of dry CH_2Cl_2 was added dropwise in 20 mins. The solution was stirred 30 mins in a cooling bath and for 2 h at room temperature. The mixture was poured on ice and 10 mL of concentrated HCl and stirred for 15 mins. The organic layer was separated, washed with H_2O and saturated brine, and dried. The solvent was removed and the residue chromatographed on Florisil™. Hexane-ether (9:1) eluted 3.37 g of the oily title compound.

1-[2-(9-(or 10)-Methyl-9,10-dihydrophenanthryl)]-1-ethanol (11)

To a stirred solution of 980 mg of 2-acetyl-9-(or 10)-methyl-9,10-dihydrophenanthrene (10) in 40 mL of MeOH, 1 g of NaBH₄ was added in small portions within 45 mins and stirring was continued for 2 h more. About 30 mL of MeOH was distilled off under reduced pressure. The residue was taken up in CH₂Cl₂; the solution was washed with H₂O and dried; and the solvent was removed. The title compound (944 mg) was obtained as an oil.

2-Acetoxyphenanthrene (12)

A solution of 2.2 g (10 mmol) of 2-acetylphenanthrene in 50 mL of CH₂Cl₂ was stirred with 3.45 g (20 mmol) of m-chloro-peroxybenzoic acid for 3 h and allowed to stand for 93 h at room temperature. The solution was washed twice with 5% KOH solution and once with H₂O, was dried and the solvent removed. From MeOH there was obtained 1.52 g of the title compound, m.p. 141-142.5°.

2-Hydroxy-9,10-dihydrophenanthrene (13)

One gram of 2-acetoxyphenanthrene was hydrogenated in 30 mL of ethyl acetate with 0.5 g of 10% Pd/C catalyst for 120 h at 35 lb/sq.i. pressure. The catalyst was filtered off and washed with ethyl acetate. The filtrate was evaporated to dryness.

The product obtained from the hydrogenation was dissolved in 50 mL of MeOH, 1 mL of concentrated HCl was added and the solution was allowed to stand for 23 h at room temperature. Two grams of NaOAc was added, the NaCl filtered off and most of the solvent was removed from the filtrate. The residue was dissolved in EtOAc, the solution washed with NaHCO₃ solution, dried, and the solvent was removed under reduced pressure. The residue (943 mg) was chromatographed on Florisil™. Hexanebenzene (6:4) eluted 480 mg of the oily 2-hydroxy-9,10-dihydrophenanthrene.

Shown in Figure 3 is a schematic representation of the synthesis of compounds 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, and 24, as described below. Also described below is the synthesis of derivative 25.

Ethyl 1-Hydroxy-5-methoxy-1,2,3,4-tetrahydronaphthalene-1-acetate (14)

To 300 mL of 1 M lithium bis-(trimethylsilyl)amide in THF, 26.43 g (0.3 mole) of dry ethyl acetate was added dropwise at -78°C under N₂. The solution was stirred for 15 mins, then a solution of 52.87 g (0.3 mole) of 5-methoxy-1-tetralone in 200 mL of dry THF was added over 40 mins. Stirring was continued for 10 mins more, then 75 mL of 20% HCl was added over 10 mins. The mixture was allowed to come to room temperature, then the organic layer was diluted with benzene, and washed with H₂O. After drying, the solvents were removed under reduced pressure. The residue crystallized upon addition of hexane to yield the title compound, 62.17 g (m.p. 57-58°C), second crop, 12.80 g (m.p. 50-53°C) yield 94.4%.

Ethyl 5-Methoxynaphthalene-1-acetate (15)

The above ester (14) 74.9 g) was heated at reflux in 60 mL of benzene with 600 mg of TsOH for 30 mins. The solution was cooled, washed with NaHCO₃ solution, and dried. Evaporation of the solvent left 69.4 g (99.5%) of an oily mixture of products with the newly formed double bond in the endocyclic and exocyclic positions.

A solution of the above product in 1400 mL of triethylene glycol dimethyl ether was heated at reflux with 30 g of 10% palladium-carbon catalyst under N₂ for 2 h. The mixture was cooled, the catalyst filtered off, and the filtrate was diluted with 3600 mL of ice-water and allowed to stand in the cold overnight. The crystals were filtered off, washed with H₂O, and dried. After stirring with 60 mL of EtOH, the colorless crystals were filtered off and dried to yield the title compound (47.4 g, 68.3%) melting at 62.5-65.6°C, m.p. of a recrystallized sample 67.5-68.5°C.

1-(2-Hydroxyethyl)-5-methoxynaphthalene (16)

To a stirred suspension of 3 g (0.08 mole) of LiAlH₄ in 100 mL of dry Et₂O, a solution of 30 g (0.12 mole) of ethyl-5-methoxy-naphthalene-1-acetate in 400 mL of dry Et₂O was added over 135 mins under N₂. The mixture was stirred for 2 h, cooled in ice, and decomposed by dropwise addition of 20 mL of satd. aq. Na₂SO₄ solution. The precipitate was filtered off and washed with Et₂O. After evaporation of the solvent a

crystalline product remained that was stirred with cold hexane. The alcohol (21.95 g, 90.5%) melted at 61-62 °C.

1-(2-Iodoethyl)-5-methoxynaphthalene (17)

To a solution of 18.4 g (0.032 mole) of P_2I_4 in 600 mL of dry CS_2 , a solution of 21.9 g (0.108 mole) of the above alcohol (16) in 250 mL of CS_2 was added at 0 °C under N_2 . The dark solution was stirred at room temperature for 120 h. Solid K_2CO_3 (25 g) and 140 mL of satd. K_2CO_3 solution were added and the mixture was stirred for 20 mins. The organic layer was separated, dried, and the solvent removed. The residue was absorbed on 125 g of Florisil™. The column was eluted with 1200 mL of hexane to yield the crystalline iodo compound (24.80 g, 73.6%) melting at 45-47 °C.

2-[2-(5-Methoxy-1-naphthyl)ethyl] cyclohexan-1,4-dione (18)

The solution of 48.9 mL (88 mmol) of 1.8 M t-BuLi in 450 mL of dry THF, cooled to -78 °C, and 12.34 g (88 mmol) of 1,4-dimethoxycyclohexa-1,4-diene was added under N_2 . After stirring for 1 h, 16 mL (92 mmol) of hexamethylphosphoramide (HMPA) was added and the deep red solution was stirred for 10 mins. The iodo compound (24 g, 77 mmol) (17) in 75 mL of dry THF was then added over 5 mins. After stirring 10 more mins, the solution was decomposed with 150 mL of saturated brine and extracted twice with hexane. The extracts were washed twice with 75 mL portions of brine, combined, dried and the solvents were removed under reduced pressure at 40 °C bath temperature.

The residue (25.23 g) was dissolved in 500 mL of acetone, the solution purged with N_2 for 20 mins and 167 mL of 1 N HCl, purged previously with N_2 , was added to the vigorously stirred solution. Stirring was continued for 1 h. The acetone was removed under reduced pressure and the cooled residue was extracted twice with CH_2Cl_2 . The extracts were washed with H_2O dried, and the solvent was removed. The residue was stirred with warm MeOH, the insoluble material (1.39 g, m.p. 179-184 °C) was filtered off and the filtrate evaporated to dryness. The residue was crystallized from Et₂O to yield the title compound (15.61 g, 88.4%), m.p. 94-100 °C. A recrystallized sample melted at 100-101.5 °C.

1-Methoxy-5,6,10,10a-tetrahydro-8(9H)benzo[a]phenanthrene-8-one (19)

To a stirred solution of 15 g of the above diketone (18), in 1500 mL of CH_2Cl_2 , 100 mL (151 g) of $MeSO_3H$ was added over 50 mins under N_2 . Stirring was continued for 10 h more. The solution was poured on ice, and the organic layer was washed twice with H_2O and dried. The residue was stirred with 40 mL of benzene. The α,β -unsaturated ketone (9.50 g, 67.5%) had a melting point of 172-177.5 °C. A recrystallized sample (acetone) melted at 177-179 °C.

Ethylene ketal of 1-methoxy-4'-oxo-7,8-cyclohexeno-phenanthrene (20)

The above ketone (38.2 g) (19) in 1300 mL of benzene was heated at reflux with 2.6 g of TsOH and 55 mL of ethylene glycol for 21 h using a Dean-Stark trap. After cooling, the solution was washed with $NaHCO_3$ solution, dried, and the solvent removed to give 44.06 g (100%) of the ethylene ketal of 1-methoxy-5,6,10,10a-tetrahydro-8(9H)benzo[a]phenanthrene-8-one (19) which was dissolved in 1000 mL of dry benzene and heated at reflux with 36 g (1.1 equiv.) of DDQ for 15 mins under N_2 . After cooling, the hydroquinone was filtered off and washed with benzene. The filtrate was concentrated to 500 mL and filtered through a column of 350 g of Florisil™. The column was eluted with a total of 4 L of benzene to yield the dehydrogenated compound (39.1 g, 89.4%), m.p. 162-163 °C (from acetone with a few drops of pyridine).

Ethylene ketal of 1-methoxy-4'-oxo-7,8-cyclohexeno-9,10-dihydrophenanthrene (21)

To 350 mL of anhydrous liquid NH_3 and 300 mg of dry $FeCl_3$, a solution of 10.00 g (31.2 mmol) of the above tetrahydro ketal (20) in 400 mL of dry THF was added with stirring under He. Li wire (1.080 g; 155.8 mg-atom; 5 equiv.) was added in small pieces. The ammonia was allowed to reflux for 1 h. The reaction was quenched with NH_4Cl and the mixture poured on ice. The mixture was extracted twice with CH_2Cl_2 , the extracts washed twice with H_2O , combined, dried, filtered through Celite and the solvent was removed under reduced pressure.

The residue, 9.06 g, in 250 mL of benzene was heated at reflux with 10 mL of ethylene glycol and 500 mg of TsOH for 22 h under N₂ using a Dean-Stark water trap. After cooling, the solution was washed with NaHCO₃, dried, reduced to a small volume and adsorbed on 100 g of Florisil™. The column was eluted with benzene. The eluted material was crystallized from Et₂O to yield (5.89 g, 58%) of the compound (21) m.p. 109-110 °C.

4'-oxo-7,8-cyclohexeno,9,10-dihydrophenanthren-1-ol (22)

To a stirred solution of 5.59 g (90 mmol) of EtSH in 40 mL of dry DMF, cooled in ice, 2.16 g (3.60 g of 60% oil dispersion; 90 mmol) of NaH was added under N₂. After the vigorous reaction subsided, the mixture was stirred at room temperature for 15 mins. A solution of 2.90 g (9 mmol) of the above ethylene ketal (21) in 40 mL of dry DMF was added, and the mixture was heated at reflux for 1 h. The mixture was poured into ice-water and extracted twice with EtOAc; the extracts were washed twice with H₂O, dried, and the solvent was removed. The residue was adsorbed on 30 g of Florisil™. The column was washed with hexane to remove the oil and then eluted with benzene-Et₂O (85:15) to give 2.36 g of the demethylated product. The product was dissolved in 40 mL of acetone, the solution purged with N₂, 400 mg of TsOH was added, and the solution stirred for 4 h under N₂. Part of the ketone separated during the reaction. Water (80 mL) was slowly added, and the crystals were filtered off, washed with acetone H₂O, and dried to give 1.92 g (80.7%) of the free ketone, m.p. 242-247 °C.

4'-oxo-7,8-cyclohexeno,9,10-dihydrophenanthren-1-ol Propionate (23)

Sixty-five mg of 4'-oxo-7,8-cyclohexeno,9,10-dihydrophenanthren-1-ol (22) was dissolved in 2 mL of dry pyridine and 1 mL of propionic anhydride. The solution was allowed to stand 20 h at room temperature and was then poured into ice water. The mixture was extracted with benzene, the extract was washed with 1 N HCl and 5% NaHCO₃ solution and dried. The colored solution was filtered through a short column of Florisil™ and the filtrate evaporated to dryness to give 62 mg of crystalline propionate.

4'-oxo-7,8-cyclohexeno,9,10-dihydrophenanthren-1-ol Acetate (24)

Fifty mg of 4'-oxo-7,8-cyclohexeno,9,10-dihydrophenanthren-1-ol (22) in 1.5 mL of dry pyridine and 0.5 mL of acetic anhydride was set aside for 24 h at room temperature and was then poured into ice water. The mixture was extracted with benzene, the extract was washed with 1 N HCl and 5% NaHCO₃ solution, dried and filtered through a short column of Florisil™. Evaporation of the solvent under reduced pressure left 50 mg of crystalline acetate.

4'-oxo-7,8-cyclohexenophenanthren-1-ol (25)

To a stirred solution of 621 mg (10 mmol) of EtSH in 5 mL of anh., dimethylformamide, cooled in ice, 240 mg (10 mmol) of NaH was added under N₂. The mixture was stirred at room temperature for 15 mins. A solution of 320 mg (1 mmol) of compound (20) in 5 mL of anh. dimethylformamide was added and heated at reflux for 1 h. The mixture was poured on ice, extracted with benzene, the extract was washed twice with H₂O, dried, and the solvent removed. The residue was adsorbed on Florisil™. Benzene-ether (25:15) eluted 285 mg of material which was dissolved in 8 mL of acetone. Forty mg of tosic acid was added and the solution stirred for 4 h at room temperature. Water was added and the precipitated produce (25) was filtered off, washed with acetone-H₂O (1:1) and dried.

Shown in Figure 4 is a schematic representation of the synthesis of compounds 26, 27, 28, and 29 as described below.

2-Bromoethyl-9,10-dihydrophenanthrene (28)

To a stirred solution of 5.56 (25 mmol) of (8) in 100 mL of THF, 9.78 g (26 mmol) of phenyltrimethylammonium tribromide was added. After stirring for 20 min, the solution became pale yellow and a precipitate of phenyltrimethylammonium bromide separated. Water (300 mL) was added and the mixture was extracted with ethyl acetate. The extract was washed with water and evaporated to dryness to yield (28) which was further purified by titration with ether and recrystallized from methanol to furnish pure (28), 5.66 g; m.p. 82-83 °C.

2-Oxiranyl,9,10-dihydrophenanthrene (29)

To a boiling solution of 1.51 g (5 mmol) of (28) in ethanol (40 mL), a solution of 285 mg (7.5 mmol) of NaBH₄ in 5 mL of water was added dropwise. Boiling was continued for 5 min, the solution was cooled, diluted with water, and extracted with ether. The ether extracted was washed with water, dried, and the solvent was removed at reduced pressure to provide (29) as a colorless oil. This solidified on trituration with a few drops of ether to yield (29), m.p. 65-66 °C.

2-[2-(9,10-dihydrophenanthryl)]-1-ethanol (26)

A solution of (29) (778 mg, 3.5 mmol) in 10 mL of dry THF was added dropwise to a stirred suspension of 152 mg (4 mmol) of lithium aluminium hydride in 10 mL of dry THF under nitrogen. The reaction mixture was stirred for 19 h. Saturated sodium sulfate solution was added dropwise to the mixture with cooling, and the precipitate was filtered off, and washed with THF. The filtrate was evaporated to dryness, and the product was crystallized from ether-hexane to yield (26), 587 mg; m.p. 68-70 °C.

1',2'-Dihydroxyethyl-9,10-dihydrophenanthrene (27)

A solution of 500 mg of (19) in 15 mL of acetone and 1.5 mL of water was stirred with p-toluenesulfonic acid (165 mg) for 5 h. An additional 330 mg of the acid was added and stirring was continued for 23 h. Then an additional 400 mg of the acid and 1.5 mL of water was added, and stirring was continued for 75 h. The mixture was extracted with ether, the extract was washed with sodium bicarbonate solution, dried, and the solvent was removed to provide crude (27), 509 mg. This was taken up in pyridine (10 mL), treated with acetic anhydride, and the solution was allowed to stand for 21 h at room temperature. The solution was poured into ice-water, the mixture was extracted with ether, washed with 1N HCl and sodium bicarbonate solution, dried, and evaporated to dryness. The diacetate product (570 mg) was dissolved in boiling hexane and chromatographed on a column of Florisil™. Elution with benzene gave the diacetate. To a solution of the latter (476 mg, 1.47 mmol) in 20 mL of methanol was added a solution of NaOH (235 mg) in 1.5 mL of water and 2 mL of methanol. After standing at room temperature for 16.5 h, this solution was made slightly acidic with acetic acid and worked up conventionally to yield (27), 308 mg. Recrystallization from acetone-hexane gave (27), 233 mg, m.p. 132-133 °C.

EXAMPLE 2

Various methods for measuring receptor binding of radioactive steroid hormones have been described and discussed in detail in a variety of publications including Fang, S., et al., J. Biol. Chem., 248:16 (1971); Liao, S., et al., J. Biol. Chem., 248:6154 (1973); Schilling, et al., The Prostate, 5:58T (1984); Liao, S., et al., Proc. Nat'l. Acad. Sci. (USA), 82:8345 (1985). The relative androgen receptor binding activities of some of the compounds of Example 1 were tested. Both the hydroxylapatite-filter assay and the tissue incubation assay were used to measure the ability to compete with radioactive steroid hormones for binding to androgen receptors as described at page 124 of Chang, C., et al., J. Steroid Biochem., 27:123 (1987). The results are presented below in Table 1. The relative binding activity (RBA) is given by:

$$RBA = \frac{IC_{50} \text{ for R1881}}{IC_{50} \text{ for test compound}} \times 100$$

Table 1: Relative receptor binding activity (RBA). The effects of test compounds on [^3H]R1881 (17 α -methyltrienolone) binding to androgen receptor of rat ventral prostate in the cell-free assay system. The concentrations (mM) of test compounds needed to show 50% inhibition (IC_{50}) are shown. RBA values are shown in parentheses.
















Compound	$\text{IC}_{50}(\text{mM})$	Compound	$\text{IC}_{50}(\text{mM})$
 R1881	0.02 (100.00)	 17β-estradiol	1.5 (1.33)
 9,10-dihydrophenanthrene	10.0 (0.20)	 10,11-dihydro-5H-dibenzo- [a,d]cycloheptene	60.0 (0.03)
 1,2,3,4,5,6,7,8-octahydro- phenanthrene	200 (0.01)	 phenanthrene	800 (0.0025)
 4'-oxo-7,8-cyclohexeno- 9,10-dihydrophenanthren- 1-ol (22)	1.3 (1.53)	 2-hydroxy-9,10-dihydro- phenanthrene (13)	7.9 (0.25)

Table 1: Continued.

Compound	IC ₅₀ (mM)	Compound	IC ₅₀ (mM)
 4'-oxo-7,8-cyclohexeno-phenanthren-1-ol (25)	>100 (<0.02)	 (±)-2-(1'-hydroxyethyl)-9,10-dihydrophenanthrene (9)	2.6 (0.77)
 (±)-6,7,16,17-tetrahydro-15H-cyclo-penta[a]phenanthren-17-ol (7)	4.2 (0.48)	 (±)-2-(1'-hydroxyethyl)-9-(or 10)-methyl-9,10-dihydrophenanthrene (11)	6.8 (0.29)
 (±)-6,7,16,17-tetrahydro-15H-cyclo-penta[a]phenanthren-17-one (6)	17.8 (0.11)	 2-(2'-hydroxyethyl)-9,10-dihydrophenanthrene (26)	3.6 (0.56)
		 2-(2',3'-dihydroxyethyl)-9,10-dihydrophenanthrene (27)	47.9 (0.04)

The IC₅₀ is that concentration needed to give 50% inhibition of ³H-R1818 binding to androgen receptors isolated, as described in Chang, C., et al., J. Steroid Biochem., 27:123 (1987), from rat ventral prostate. Table 1 gives the IC₅₀ for reference compounds, including: R1881; 17- β -estradiol; 9,10-dihydrophenanthrene; 10,11-dihydro-5H-dibenzo[a,d]cyclopentene and for compounds of the invention including: 4'-oxo-7,8-cyclohexeno-9,10-dihydrophenanthren-1-ol (22); 1-[2-(9,10-dihydrophenanthryl)]-1-ethanol (9); 6,7,16,17-

tetrahydro-15H-cyclopenta[a]phenanthren-17-ol (7); 1-[2-(9 or 10)-methyl-9,10-dihydrophenanthryl]-1-ethanol (11); 2-hydroxy-9,10-dihydrophenanthrene (13); 6,7,16,17-tetrahydro-15H-cyclopenta[a]phenanthren-17-one (6); 4'-oxo-7,8-cyclohexenophenanthren-1-ol (25); 2-(2'-hydroxyethyl)-9,10-dihydrophenanthrene (26); and 2-(2',3'-dihydroxyethyl)-9,10-dihydrophenanthrene (27).

- 5 An IC_{50} of 10 μ M or less is considered most desirable for an effective anti-androgen. The results indicate that the generalized structure (as depicted as page 3) for the most effective anti-androgens are those derivatives which have a flexible structure between, preferably, two aromatic rings. A carbonyl group and/or a hydroxyl group at either end of the compound, as in the case with natural androgens, increases the potency, apparently by increasing the recognition of functional groups in the binding cavity of the androgen receptor.

10 Because of the possibility that the compounds of the invention may display particularly enhanced activity in certain organs, the assay methods described above, while useful in screening potential anti-androgen candidates, may not show the extent of the relative anti-androgenic potencies of the compounds in organs with different pathogenic conditions. Accordingly, it is anticipated that some of the new compounds, that may not exhibit high activities in the assays shown, may still be useful in the treatment of abnormalities. One way to additionally confirm the effectiveness of the anti-androgens of the present invention is to perform an *in vivo* anti-androgenic activity assay. Chang, C., et al., *J. Steroid Biochem.*, 27:123 (1987) (legend of Table 2, page 129) (see also "androgenicity assay" on page 6155 of Liao, S., et al., *J. of Biol. Chem.*, 248:6154 (1973). Briefly, the method comprises castrating rats on day one and injecting, with either testosterone propionate and/or a test compound, subcutaneously daily until the 8th day. On the 9th day, the rats are killed and organs, such as the ventral prostate, the seminal vesicle, and the coagulating gland, from individual rats are weighed.

Excessive androgenic activities are implicated in pathogenesis of certain androgen dependent conditions, such as benign prostatic hyperplasia and prostate cancer (Huggins, et al., *Cancer Res.*, 1:293 (1941); Silteri, et al., *J. Clin. Invest.*, 49:1737 (1970); Geller, et al., *J. Clin. Endocr. Metab.*, 43:686 (1976); acne (Sansone, et al., *J. Invest. Dermat.*, 56:366 (1971)), male pattern baldness (Bingham, et al., *J. Endocr.*, 59:11 (1973); and female hirsutism (Kutten, et al., *J. Endocr.*, 25:83 (1977)). The anti-androgens of the invention, are therefore expected to be useful in treating these abnormalities [(Martini, L. and Motta, M., *Androgens and Anti-androgens*, Raven Press, New York (1977)) because many of those abnormalities are due to excess androgens and androgenic activities in the organs affected.

Compounds (active or inactive in the assays shown) can be administered by topical (especially for skin) or internal (oral or injection) routes. In addition, it is expected that some of these compounds may interact with mutated receptors and, therefore, may be useful in the treatment, as well as in the diagnosis of androgen and other hormone-insensitive tumors. The compounds of the invention are also expected to be important in the studies of the mechanism of action of androgens and anti-androgens especially if they are radioactively labelled.

Further, it is anticipated that the compounds of the invention will find utility in localizing androgen receptors. For example, compounds of the invention can be suitably labelled (e.g., with fluorescent groups or radioactive or non-radioactive isotopes) to serve as markers for both *in vitro* and *in vivo* analyses of, for example, the tissue distribution of androgen receptors.


It is envisioned that various dihydrophenanthrene derivatives and compounds with steroid ring systems, as described above, will also be effective according to the present invention. Although the preferred compounds are: 6,7,16,17-tetrahydro-15H cyclopenta[a] phenanthren-17-ol (7); 1-[2-(9,10-dihydrophenanthryl)-1-ethanol (9); 1-[2-(9 or 10)-methyl-9,10-dihydrophenanthryl]-1-ethanol (11); 2-hydroxy-9,10-dihydrophenanthrene (13); 4'-oxo-7,8-cyclohexeno,9,10-dihydrophenanthren-1-ol (22); 4'-oxo-7,8-cyclohexenophenanthren-1-ol (25); 2-(2'-hydroxyethyl)-9,10-dihydrophenanthrene (26); and 2-(2',3'-dihydroxyethyl)-9,10-dihydrophenanthrene (27); it is not intended to preclude others, such as 16,17-dihydro-15H-cyclopenta[a]phenanthrene (4); 6,7,16,17-tetrahydro-15H-cyclopenta[a]phenanthrene (5); 2-acetyl-9,10-dihydrophenanthrene (8); 2-acetyl-9(or 10)-methyl-9,10-dihydrophenanthrene (10); 2-acetoxypheanthrene (12); 1-methoxy-5,6,10a-tetrahydro-8(9H)benzo[a]phenanthrene-8-one (19); 4'-oxo-7,8-cyclohexeno,9,10-dihydrophenanthren-1-ol propionate (23); and 4'-oxo-7,8-cyclohexeno,9,10-dihydrophenanthren-1-ol acetate (24) from being included in the scope of the present invention.

Also, because the acetate and propionate forms of androgens and anti-androgens have been used effectively *in vivo*, it is expected that these esters (as well as other ester analogues) are more soluble and metabolically stable and may be more appropriate than the free-alcohol forms for the delivery to the target sites; therefore, these derivatives may be more potent when used as drugs for treating abnormalities and are also contemplated within the scope of the invention.

Claims

Claims for the following Contracting States : AT, BE, CH, DE, FR, GB, GR, IT, LI, LU, NL, SE

1. An anti-androgen comprising a compound selected from the group consisting of compounds of the general formula:




wherein said ring A is substituted or unsubstituted and wherein said ring A is substituted it is fused at either the 1-2 or 2-3 positions to a hydroxycycloalkyl, or a ketocycloalkyl, or is substituted with a hydroxyl, an alkyl, or a hydroxyalkyl other than hydroxyisopropyl other than hydroxyisopropyl at the 1, 2 or 3 positions;

wherein said ring B is saturated or mono-unsaturated at the 9-10 position and is optionally substituted with a lower alkyl at either the 9 or 10 position;

wherein said ring C is substituted or unsubstituted, and wherein when said ring C is substituted it is fused at the 6-7 or 7-8 positions to a ketocycloalkyl or a hydroxycycloalkyl; and

provided that rings A and C may not both be unsubstituted, wherein the compound is not 6,7,16,17-tetrahydro-15H-cyclopenta(a)phenanthren-17-one or 16, 17-dihydro-15H-cyclopenta(a)phenanthren-17-one, or a derivative of said compound wherein a free hydroxyl group has been replaced with an acetate or a propionate moiety.

2. The anti-androgen according to Claim 1 wherein:
said ketocycloalkyl is cyclopentanone or cyclohexanone.
3. The anti-androgen according to Claim 1 wherein:
said hydroxycycloalkyl is cyclopentanol or cyclohexanol.
4. The anti-androgen according to Claim 1, wherein:
said alkyl is methyl.
5. The anti-androgen according to Claim 1 which comprises a member selected from the group consisting of:
1-[2-(9,10-dihydrophenanthryl)]-1-ethanol; 6,7,16,17-tetrahydro-15H-cyclopenta[a]phenanthren-17-ol; 1-[2-(9(or 10)-methyl-9,10-dihydrophenanthryl)]-1-ethanol; 2-hydroxy-9,10-dihydrophenanthrene; 4'-oxo-7,8-cyclohexeno,9,10-dihydrophenanthren-1-ol; 4'-oxo 7,8-cyclohexenophenanthren-1-ol; 2-(2'-hydroxyethyl)-9,10-dihydrophenanthrene; and 2-(2',3'-dihydroxyethyl)-9,10-dihydrophenanthrene.
6. The anti-androgen according to Claim 1, which comprises a member selected from the group consisting of:
4'-oxo-7,8-cyclohexano,9,10-dihydrophenanthren-1-ol propionate; and
4'-oxo-7,8-cyclohexano,9,10-dihydrophenanthren-1-ol acetate.
7. A pharmaceutical composition for use in the treatment of disorders associated with excessive androgenic activity in an organism, said composition comprising a pharmaceutically acceptable solvent, diluent, adjuvant or carrier and, as the active ingredient, an anti-androgen comprising a compound selected from the group consisting of compounds of the general formula:




wherein said ring A is substituted or unsubstituted and wherein when said ring A is substituted it is fused at either the 1-2 or 2-3 positions to a hydroxycycloalkyl, or a ketocycloalkyl, or is substituted with a hydroxyl, an alkyl, or a hydroxyalkyl other than hydroxyisopropyl at the 1, 2, or 3 positions;

wherein said ring B is saturated or monounsaturated at the 9-10 position and is optionally substituted with a lower alkyl at either the 9 or 10 position;

wherein said ring C is substituted or unsubstituted, and wherein when said ring C is substituted it is fused at the 6-7 or 7-8 positions to a ketocycloalkyl or a hydroxycycloalkyl; and provided that rings A and C may not both be unsubstituted.

8. Use of an anti-androgen comprising a compound selected from the group consisting of compounds of the general formula:



wherein said ring A is substituted or unsubstituted and wherein when said ring A is substituted it is fused at either the 1-2 or 2-3 positions to a hydroxycycloalkyl, or a ketocycloalkyl, or is substituted with a hydroxyl, an alkyl, or a hydroxyalkyl at the 1, 2 or 3 positions;

wherein said ring B is saturated or monounsaturated at the 9-10 position and is optionally substituted with a lower alkyl at either the 9 or 10 position;

wherein said ring C is substituted or unsubstituted, and wherein when said ring C is substituted it is fused at the 6-7 or 7-8 positions to a ketocycloalkyl or a hydroxycycloalkyl; and

provided that rings A and C may not both be unsubstituted, for the manufacture of a medicament for the treatment of disorders associated with excessive androgenic activity in an organism.

9. Use according to Claim 8 wherein said anti-androgen is selected from the group consisting of:

1-[2-(9,10-dihydrophenanthryl)]-1-ethanol; 6,7,16,17-tetrahydro-15H-cyclopenta[a]phenanthren-17-ol; 1-[2-(9 or 10)-methyl-9,10-dihydrophenanthryl]-1-ethanol; 2-hydroxy-9,10-dihydrophenanthrene, 6,7,16,17-tetrahydro-15H-cyclopenta[a]phenanthren-17-one; 4'-oxo-7,8-cyclohexeno, 9,10-dihydrophenanthren-1-ol; 4'-oxo 7,8-cyclohexenophenanthren-1-ol; 2-(2'-hydroxyethyl)-9,10-dihydrophenanthrene; and 2-(2',3'-dihydroxyethyl)-9,10-dihydrophenanthrene.


10. A method for localizing androgen receptors in a sample, said method comprising the steps of:

incubating said sample with a compound according to Claim 1, wherein said compound is labelled with a marker;

detecting said marker; and
localizing said androgen receptor in said sample.

Claims for the following Contracting State : ES

1. A method for localizing androgen receptors in a sample said method comprising incubating said sample with a compound selected from the group consisting of compounds of the general formula:




wherein said ring A is substituted or unsubstituted and wherein said ring A is substituted it is fused at either the 1-2 or 2-3 positions to a hydroxycycloalkyl, or a ketocycloalkyl, or is substituted with a hydroxyl, an alkyl, or a hydroxyalkyl other than hydroxyisopropyl other than hydroxyisopropyl at the 1, 2 or 3 positions;

wherein said ring B is saturated or mono-unsaturated at the 9-10 position and is optionally substituted with a lower alkyl at either the 9 or 10 position;

wherein said ring C is substituted or unsubstituted, and wherein when said ring C is substituted it is fused at the 6-7 or 7-8 positions to a ketocycloalkyl or a hydroxycycloalkyl; and

provided that rings A and C may not both be unsubstituted, wherein the compound is not 6,7,16,17-tetrahydro-15H-cyclopenta(a)phenanthren-17-one or 16, 17-dihydro-15H-cyclopenta(a)phenanthren-17-one, or a derivative of said compound wherein a free hydroxyl group has been replaced with an acetate or a propionate moiety, wherein said compound is labelled with a marker, detecting said marker and localizing said androgen receptor in said sample.

2. The method according to Claim 1 wherein:
said ketocycloalkyl is cyclopentanone or cyclohexanone.
3. The method according to Claim 1 wherein:
said hydroxycycloalkyl is cyclopentanol or cyclohexanol.
4. The method according to Claim 1, wherein:
said alkyl is methyl.
5. The method according to Claim 1 which comprises a member selected from the group consisting of:
1-[2-(9,10-dihydrophenanthryl)]-1-ethanol; 6,7,16,17-tetrahydro-15H-cyclopenta[a]phenanthren-17-ol; 1-[2-(9(or 10)-methyl-9,10-dihydrophenanthryl)]-1-ethanol; 2-hydroxy-9,10-dihydrophenanthrene; 4'-oxo-7,8-cyclohexeno,9,10-dihydrophenanthren-1-ol; 4'-oxo 7,8-cyclohexenophenanthren-1-ol; 2-(2'-hydroxyethyl)-9,10-dihydrophenanthrene; and 2-(2',3'-dihydroxyethyl)-9,10-dihydrophenanthrene.
6. The method according to Claim 1, which comprises a member selected from the group consisting of:
4'-oxo-7,8-cyclohexano,9,10-dihydrophenanthren-1-ol propionate; and
4'-oxo-7,8-cyclohexano,9,10-dihydrophenanthren-1-ol acetate.
7. Use of an anti-androgen comprising a compound selected from the group consisting of compounds of the general formula:



5

10

15

wherein said ring A is substituted or unsubstituted and wherein when said ring A is substituted it is fused at either the 1-2 or 2-3 positions to a hydroxycycloalkyl, or a ketocycloalkyl, or is substituted with a hydroxyl, an alkyl, or a hydroxyalkyl at the 1, 2 or 3 positions;

wherein said ring B is saturated or monounsaturated at the 9-10 position and is optionally substituted with a lower alkyl at either the 9 or 10 position;

20

wherein said ring C is substituted or unsubstituted, and wherein when said ring C is substituted it is fused at the 6-7 or 7-8 positions to a ketocycloalkyl or a hydroxycycloalkyl; and

provided that rings A and C may not both be unsubstituted, for the manufacture of a medicament for the treatment of disorders associated with excessive androgenic activity in an organism.

25

8. Use according to Claim 7 wherein said anti-androgen is selected from the group consisting of:

1-[2-(9,10-dihydrophenanthryl)]-1-ethanol; 6,7,16,17-tetrahydro-15H-cyclopenta[a]phenanthren-17-ol; 1-[2-(9(or 10)-methyl-9,10-dihydrophenanthryl)]-1-ethanol; 2-hydroxy-9,10-dihydrophenanthrene; 6,7,16,17-tetrahydro-15H-cyclopenta[a]phenanthren-17-one; 4'-oxo-7,8-cyclohexeno-9,10-dihydrophenanthren-1-ol; 4'-oxo 7,8-cyclohexenophenanthren-1-ol; 2-(2'-hydroxyethyl)-9,10-dihydrophenanthrene; and 2-(2',3'-dihydroxyethyl)-9,10-dihydrophenanthrene.


30

Patentansprüche

Patentansprüche für folgende Vertragsstaaten : AT, BE, CH, DE, FR, GB, GR, IT, LI, LU, NL, SE

35

1. Anti-Androgen, das eine Verbindung umfaßt, die ausgewählt ist aus der Gruppe, die aus Verbindungen der allgemeinen Formel besteht:



40

45


50

55

wobei besagter Ring A substituiert oder nicht-substituiert ist und wobei, wenn besagter Ring A substituiert ist, er an entweder den 1-2- oder 2-3-Positionen an ein Hydroxycycloalkyl oder ein Ketocycloalkyl kondensiert ist oder mit einem Hydroxyl, einem Alkyl oder einem anderen Hydroxyalkyl als Hydroxyisopropyl an den 1-, 2- oder 3-Positionen substituiert ist; wobei besagter Ring B gesättigt oder einfach ungesättigt an der 9-10-Position ist und fakultativ mit einem niederen Alkyl an entweder der 9- oder 10-Position substituiert ist; wobei besagter Ring C substituiert oder nicht-substituiert ist und wobei, wenn besagter Ring C


substituiert ist, er an den 6-7- oder 7-8-Positionen an ein Ketocycloalkyl oder ein Hydroxycycloalkyl kondensiert ist; und vorausgesetzt die Ringe A und C können nicht beide nichtsubstituiert sein, wobei die Verbindung nicht 6,7,16,17-Tetrahydro-15H-cyclopenta(a)phenanthen-17-on oder 16,17-Dihydro-15H-cyclopenta(a)-phenanthen-17-on ist, oder ein Derivat besagter Verbindung, in der eine freie Hydroxylgruppe durch eine Acetat- oder eine Propionat-Einheit ersetzt worden ist.

2. Anti-Androgen nach Anspruch 1, wobei: besagtes Ketocycloalkyl Cyclopentanon oder Cyclohexanon ist.
3. Anti-Androgen nach Anspruch 1, wobei: besagtes Hydroxycycloalkyl Cyclopentanol oder Cyclohexanol ist.
4. Anti-Androgen nach Anspruch 1, wobei: besagtes Alkyl Methyl ist.
5. Anti-Androgen nach Anspruch 1, das ein Mitglied umfaßt, das ausgewählt ist aus der Gruppe, bestehend aus:
1-[2-(9,10-Dihydrophenantril)-1-ethanol; 6,7,16,17-Tetrahydro-15H-cyclopenta(a)phenanthen-17-ol; 1[2-(9(oder 10)-Methyl-9,10-dihydrophenantril)-1-ethanol; 2-Hydroxy-9,10-dihydrophenanthen; 4'-Oxo-7,8-cyclohexeno-9,10-dihydrophenanthen-1-ol; 4'-Oxo-7,8-cyclohexenophenanthen-1-ol; 2-(2'-Hydroxyethyl)-9,10-dihydrophenanthen; und 2-(2',3'-Dihydroxyethyl)-9,10-dihydrophenanthen.
6. Anti-Androgen nach Anspruch 1, das ein Mitglied umfaßt, das ausgewählt ist aus der Gruppe, bestehend aus:
4'-Oxo-7,8-cyclohexano-9,10-dihydrophenanthen-1-ol-propionat; und
4'-Oxo-7,8-cyclohexano-9,10-dihydrophenanthen-1-ol-acetat.
7. Pharmazeutische Zusammensetzung zur Verwendung bei der Behandlung von Erkrankungen, die mit übermäßiger androgener Aktivität in einem Organismus in Zusammenhang stehen, wobei besagte Zusammensetzung ein pharmazeutisch annehmbares Lösungsmittel, Verdünnungsmittel, Adjuvans oder Trägermittel und, als den aktiven Bestandteil, ein Anti-Androgen umfaßt, das eine Verbindung umfaßt, die ausgewählt ist aus der Gruppe, die aus Verbindungen der allgemeinen Formel besteht:



wobei besagter Ring A substituiert oder nicht-substituiert ist und wobei, wenn besagter Ring A substituiert ist, er an entweder den 1-2- oder 2-3-Positionen an ein Hydroxycycloalkyl oder ein Ketocycloalkyl kondensiert ist oder mit einem Hydroxyl, einem Alkyl oder einem anderen Hydroxyalkyl als Hydroxyisopropyl an den 1-, 2- oder 3-Positionen substituiert ist; wobei besagter Ring B gesättigt oder einfach ungesättigt an der 9-10-Position ist und fakultativ mit einem niederen Alkyl an entweder der 9- oder 10-Position substituiert ist; wobei besagter Ring C substituiert oder nicht-substituiert ist und wobei, wenn besagter Ring C substituiert ist, er an den 6-7- oder 7-8-Positionen an ein Ketocycloalkyl oder ein Hydroxycycloalkyl kondensiert ist; und vorausgesetzt die Ringe A und C können nicht beide nichtsubstituiert sein.

8. Verwendung einem Anti-Androgens, das eine Verbindung umfaßt, die ausgewählt ist aus der Gruppe, die aus Verbindungen der allgemeinen Formel besteht:



5

10

15

wobei besagter Ring A substituiert oder nicht-substituiert ist und wobei, wenn besagter Ring A substituiert ist, er an entweder den 1-2- oder 2-3-Positionen an ein Hydroxycycloalkyl oder ein Ketocycloalkyl kondensiert ist oder mit einem Hydroxyl, einen Alkyl oder einem Hydroxyalkyl an den 1-, 2- oder 3-Positionen substituiert ist;

20

wobei besagter Ring B gesättigt oder einfach ungesättigt an der 9-10-Position ist und fakultativ mit einem niederen Alkyl an entweder der 9- oder 10-Position substituiert ist;

wobei besagter Ring C substituiert oder nicht-substituiert ist und wobei, wenn besagter Ring C substituiert ist, er an den 6-7- oder 7-8-Positionen an ein Ketocycloalkyl oder ein Hydroxycycloalkyl kondensiert ist; und

25

vorausgesetzt die Ringe A und C können nicht beide nichtsubstituiert sein, für die Herstellung eines Arzneimittels für die Behandlung von Erkrankungen, die mit übermäßiger androgener Aktivität in einem Organismus in Zusammenhang stehen.

30

9. Verwendung nach Anspruch 8, wobei besagtes Anti-Androgen ausgewählt ist aus der Gruppe, bestehend aus:

1-[2-(9,10-Dihydrophenanthryl)]-1-ethanol; 6,7,16,17-Tetrahydro-15H-cyclopenta[a]phenantren-17-ol; 1[2-(9(oder 10)-Methyl-9,10-dihydrophenanthryl)]-1-ethanol; 2-Hydroxy-9,10-dihydrophenantren; 6,7,16,17-Tetrahydro-15H-cyclopenta[a]phenantren-17-on; 4'-Oxo-7,8-cyclohexeno-9,10-dihydrophenantren-1-ol; 4'-Oxo-7,8-cyclohexenophenantren-1-ol; 2-(2'-Hydroxyethyl)-9,10-dihydrophenantren; und 2-(2',3'-Dihydroxyethyl)-9,10-dihydrophenantren.

35

10. Verfahren zur Lokalisierung von Androgen-Rezeptoren in einer Probe, wobei besagtes Verfahren die Schritte umfaßt:

Inkubieren besagter Probe mit einer Verbindung gemäß Anspruch 1, wobei besagte Verbindung mit einer Markierungssubstanz markiert ist;

40

Nachweisen besagter Markierungssubstanz; und

Lokalisieren besagten Androgen-Rezeptors in besagter Probe.


45

Patentansprüche für folgenden Vertragsstaat : ES

1. Verfahren zur Lokalisierung von Androgen-Rezeptoren in einer Probe, wobei besagtes Verfahren Inkubieren besagter Probe mit einer Verbindung, die aus der Gruppe ausgewählt ist, die aus Verbindungen der allgemeinen Formel besteht:


50

55



wobei besagter Ring A substituiert oder nicht-substituiert ist und wobei, wenn besagter Ring A substituiert ist, er an entweder den 1-2- oder 2-3-Positionen an ein Hydroxycycloalkyl oder ein Ketocycloalkyl kondensiert ist oder mit einem Hydroxyl einem Alkyl oder einem anderen Hydroxyalkyl als Hydroxyisopropyl an den 1-, 2- oder 3-Positionen substituiert ist; wobei besagter Ring B gesättigt oder einfach ungesättigt an der 9-10-Position ist und fakultativ mit einem niederen Alkyl an entweder der 9- oder 10-Position substituiert ist; wobei besagter Ring C substituiert oder nicht-substituiert ist und wobei wenn besagter Ring C substituiert ist, er an den 6-7- oder 7-8-Positionen an ein Ketocycloalkyl oder ein Hydroxycycloalkyl kondensiert ist; und vorausgesetzt die Ringe A und C können nicht beide nichtsubstituiert sein, wobei die Verbindung nicht 6,7,16,17-Tetrahydro-15H-cyclopenta(a)phenantren-17-on oder 16,17-Dihydro-15H-cyclopenta(a)-phenantren-17-on ist, oder ein Derivat besagter Verbindung, in der eine freie Hydroxylgruppe durch eine Acetat- oder eine Propionat-Einheit ersetzt worden ist, wobei besagte Verbindung mit einer Markierungssubstanz markiert ist, Nachweisen besagter Markierungssubstanz und Lokalisieren besagten Androgen-Rezeptors in besagter Probe umfaßt.

2. Verfahren nach Anspruch 1, wobei: besagtes Ketocycloalkyl Cyclopentanon oder Cyclohexanon ist.
3. Verfahren nach Anspruch 1, wobei: besagtes Hydroxycycloalkyl Cyclopentanol oder Cyclohexanol ist.
4. Verfahren nach Anspruch 1, wobei: besagtes Alkyl Methyl ist.
5. Verfahren nach Anspruch 1, das ein Mitglied umfaßt, das ausgewählt ist aus der Gruppe, bestehend aus:
1-[2-(9,10-Dihydrophenantril)]-1-ethanol; 6,7,16,17-Tetrahydro-15H-cyclopenta[a]phenantren-17-ol; 1[2-(9-(oder 10)-Methyl-9,10-dihydrophenantril)]-1-ethanol; 2-Hydroxy-9,10-dihydrophenantren; 4'-Oxo-7,8-cyclohexeno-9,10-dihydrophenantren-1-ol; 4'-Oxo-7,8-cyclohexenophenantren-1-ol; 2-(2'-Hydroxyethyl)-9,10-dihydrophenantren; und 2-(2',3'-Dihydroxyethyl)-9,10-dihydrophenantren.
6. Verfahren nach Anspruch 1, das ein Mitglied umfaßt, das aus der Gruppe ausgewählt ist, bestehend aus:
4'-Oxo-7,8-cyclohexano-9,10-dihydrophenantren-1-ol-propionat; und
4'-Oxo-7,8-cyclohexano-9,10-dihydrophenantren-1-ol-acetat.
7. Verwendung eines Anti-Androgens, das eine Verbindung umfaßt, die aus der Gruppe ausgewählt ist, die aus Verbindungen der allgemeinen Formel besteht:



wobei besagter Ring A substituiert oder nicht-substituiert ist und wobei, wenn besagter Ring A substituiert ist, er an entweder den 1-2- oder 2-3-Positionen an ein Hydroxycycloalkyl oder ein Ketocycloalkyl kondensiert ist oder mit einem Hydroxyl, einem Alkyl oder einem Hydroxyalkyl an den 1-, 2- oder 3-Positionen substituiert ist;

wobei besagter Ring B gesättigt oder einfach ungesättigt an der 9-10-Position ist und fakultativ mit einem niederen Alkyl an entweder der 9- oder 10-Position substituiert ist;

wobei besagter Ring C substituiert oder nicht-substituiert ist und wobei, wenn besagter Ring C substituiert ist, er an den 6-7- oder 7-8-Positionen an ein Ketocycloalkyl oder ein Hydroxycycloalkyl kondensiert ist; und

vorausgesetzt die Ringe A und C können nicht beide nichtsubstituiert sein, für die Herstellung eines Arzneimittels für die Behandlung von Erkrankungen, die mit übermäßiger androgener Aktivität in einem Organismus in Zusammenhang stehen.


8. Verwendung nach Anspruch 7, wobei besagtes Anti-Androgen ausgewählt ist aus der Gruppe, bestehend aus;

1-[2-(9,10-Dihydrophenantryl)]-1-ethanol; 6,7,16,17-Tetrahydro-15H-cyclopenta[a]phenanthen-17-ol; 1[2-(9(oder 10)-Methyl-9,10-dihydrophenantryl)]-1-ethanol; 2-Hydroxy-9,10-dihydrophenanthen; 6,7,16,17-Tetrahydro-15H-cyclopenta[a]phenanthen-17-on; 4'-Oxo-7,8-cyclohexeno-9,10-dihydrophenanthen-1-ol; 4'-Oxo-7,8-cyclohexenophenanthren-1-ol; 2-(2'-Hydroxyethyl)-9,10-dihydrophenanthen; und 2-(2',3'-Dihydroxyethyl)-9,10-dihydrophenanthen.

Revendications

Revendications pour les Etats contractants suivants : AT, BE, CH, DE, FR, GB, GR, IT, LI, LU, NL, SE

1. Antiandrogène comprenant un composé choisi dans le groupe formé par les composés de formule générale ;




dans laquelle ledit cycle A est substitué ou non substitué et dans laquelle lorsque ledit cycle A est substitué, il est condensé en position 1-2 ou 2-3 avec un composé hydroxycycloalkylique ou céto-cycloalkylique, ou il est substitué par un groupe hydroxyle, alkyle ou un groupe hydroxyalkyle différent d'un groupe hydroxyisopropyle en position 1, 2 ou 3,

dans laquelle ledit cycle B est saturé ou mono-insaturé en position 9-10 et est éventuellement substitué par un groupe alkyle inférieur en position 9 ou en position 10,

dans laquelle ledit cycle C est substitué ou non substitué, et dans laquelle lorsque ledit cycle C est substitué, il est condensé en position 6-7 ou 7-8 avec un composé cétycycloalkylique ou hydroxycycloalkylique ; et

à condition que les cycles A et C ne puissent par être l'un et l'autre non substitués, dans lequel le composé n'est pas la 6,7,16,17-tétrahydro-15H-cyclopenta(a)phénanthrén-17-one ou la 16,17-dihydro-15H-cyclopenta(a)phénanthrén-17-one, ou un dérivé dudit composé dans lequel un groupe hydroxyle libre a été remplacé par une fraction acétate ou propionate.

2. Antiandrogène selon la revendication 1, dans lequel ledit composé cétycycloalkylique est la cyclopentanone ou la cyclohexanone.
3. Antiandrogène selon la revendication 1, dans lequel ledit composé hydroxycycloalkylique est le cyclopentanol ou le cyclohexanol.
4. antiandrogène selon la revendication 1, dans lequel ledit groupe alkyle est un groupe méthyle.
5. Antiandrogène selon la revendication 1, qui comprend un élément choisi dans le groupe formé par :
le 1-[2-(9,10-dihydrophénanthryl)]-1-éthanol, le 6,7,16,17-tétrahydro-15H-cyclopenta[a]phénanthrén-17-ol, le 1-[2-(9 ou 10)-méthyl-9, 10-dihydrophénanthryl)]-1-éthanol, le 2-hydroxy-9,10-dihydrophénanthrène, le 4'-oxo-7,8-cyclohexéno-9,10-dihydrophénanthrén-1-ol, le 4'-oxo-7,8-cyclohexénophénanthrén-1-ol, le 2-(2'-hydroxyéthyl) 9,10-dihydrophénanthrène et le 2-(2',3'-dihydroxyéthyl)-9,10-dihydrophénanthrène.
6. Antiandrogène selon la revendication 1, qui comprend un élément choisi dans le groupe formé par :
le propionate de 4'-oxo-7,8-cyclohexéno-9,10-dihydrophénanthrén-1-yle et l'acétate de 4'-oxo-7,8-cyclohexéno-9,10-dihydrophénanthrén-1-yle.
7. Composition pharmaceutique utilisable dans le traitement des troubles associés à une activité androgène excessive dans un organisme, ladite composition comprenant un solvant, diluant, adjuvant ou véhicule acceptable du point de vue pharmaceutique et, comme ingrédient actif, un antiandrogène comprenant un composé choisi dans le groupe formé par les composés de formule générale :




dans laquelle ledit cycle A est substitué ou non substitué et dans laquelle lorsque ledit cycle A est substitué, il est condensé en position 1-2 ou 2-3 avec un composé hydroxycycloalkylique ou cétycycloalkylique, ou il est substitué par un groupe hydroxyle, alkyle ou un groupe hydroxyalkyle différent d'un groupe hydroxyisopropyle en position 1, 2 ou 3,

dans laquelle ledit cycle B est saturé ou mono-insaturé en position 9-10 et est éventuellement substitué par un groupe alkyle inférieur en position 9 ou en position 10,

dans laquelle ledit cycle C est substitué ou non substitué, et dans laquelle lorsque ledit cycle C est substitué il est condensé en position 6-7 ou 7-8 avec un composé cétycycloalkylique ou hydroxycycloalkylique ; et

à condition que les cycles A et C ne puissent pas être l'un et l'autre non substitués.

8. Utilisation d'un antiandrogène comprenant un composé choisi dans le groupe formé par les composés de formule générale



dans laquelle ledit cycle A est substitué ou non substitué et dans laquelle lorsque ledit cycle A est substitué, il est condensé en position 1-2 ou 2-3 avec un composé hydroxycycloalkylique ou céto-cycloalkylique, ou il est substitué par un groupe hydroxyle, alkyle ou hydroxyalkyle en position 1, 2 ou 3,

dans laquelle ledit cycle B est saturé ou mono-insaturé en position 9-10 et est éventuellement substitué par un groupe alkyle inférieur en position 9 ou en position 10,


dans laquelle ledit cycle C est substitué ou non substitué, et dans laquelle lorsque ledit cycle C est substitué il est condensé en position 6-7 ou 7-8 avec un composé céto-cycloalkylique ou hydroxycycloalkylique ; et

à condition que les cycles A et C ne puissent pas être l'un et l'autre non substitués, pour la préparation d'un médicament pour le traitement des troubles associés à une activité androgène excessive dans un organisme.

9. Utilisation selon la revendication 8, dans laquelle ledit androgène est choisi dans le groupe formé par :
- le 1-[2-(9,10-dihydrophénanthryl)]-1-éthanol, le 6,7,16,17-tétrahydro-15H-cyclopenta[a]-phénanthrén-17-ol, le 1-[2-(9 ou 10)-méthyl-9, 10-dihydrophénanthryl)]-1-éthanol, le 2-hydroxy-9,10-dihydrophénanthrène, la 6,7,16,17-tétrahydro-15H-cyclopenta[a]phénanthrén-17-one, le 4'-oxo-7,8-cyclohexéno-9,10-dihydrophénanthrén-1-ol, le 4' oxo-7,8-cyclohexénophénanthrén-1-ol, le 2-(2'-hydroxyéthyl)-9,10-dihydrophénanthrène et le 2-(2', 3'dihydroxyéthyl)-9,10-dihydrophénanthrène,
10. Procédé pour localiser les récepteurs d'androgènes dans un échantillon ledit procédé comprenant les étapes consistant à :
- incuber ledit échantillon avec un composé selon la revendication 1, ledit composé étant marqué par un marqueur,
 - détecter ledit marqueur, et
 - localiser ledit récepteur d'androgène dans ledit échantillon.

Revendications pour l'Etat contractant suivant : ES

1. Procédé pour localiser des récepteurs d'androgènes dans un échantillon, ledit procédé comprenant l'incubation dudit échantillon avec un composé choisi dans le groupe formé par les composés de formule générale :




dans laquelle ledit cycle A est substitué ou non substitué et dans laquelle lorsque ledit cycle A est substitué, il est condensé en position 1-2 ou en position 2-3 avec un composé hydroxycycloalkylique ou un composé céto-cycloalkylique, ou il est substitué par un groupe hydroxyle, alkyle ou un groupe hydroxyalkyle différent du groupe hydroxyisopropyle en position 1, 2 ou 3,

dans laquelle ledit cycle B est saturé ou mono-insaturé en position 9-10 et est éventuellement substitué par un groupe alkyle inférieur en position 9 ou en position 10,

dans laquelle ledit cycle C est substitué ou non substitué, et dans laquelle lorsque ledit cycle C est substitué il est condensé en position 6-7 ou 7-8 avec un composé céto-cycloalkylique ou hydroxycycloalkylique ; et

à condition que les cycles A et C ne puissent pas être l'un et l'autre non substitués, dans lequel le composé n'est pas la 6,7,16,17-tétrahydro-15H-cyclopenta(a)phénanthrén-17-one ou la 16,17-dihydro-15H-cyclopenta(a)-phénanthrén-17-one, ou un dérivé dudit composé dans lequel un groupe hydroxyle libre a été remplacé par une fraction acétate ou propionate, ledit composé étant marqué par un marqueur, et pour détecter ledit marqueur et localiser ledit récepteur d'androgène dans ledit échantillon.

2. Procédé selon la revendication 1, dans lequel ledit composé céto-cycloalkylique est la cyclopentanone ou la cyclohexanone.
3. Procédé selon la revendication 1, dans lequel ledit composé hydroxycycloalkylique est le cyclopentanol ou le cyclohexanol.
4. Procédé selon revendication 1, dans lequel ledit groupe alkyle est un groupe méthyle.
5. Procédé selon la revendication 1, qui comprend un élément choisi dans le groupe formé par :
le 1-[2-(9,10-dihydrophénanthryl)]-1-éthanol, le 6,7,16,17-tétrahydro-15H-cyclopenta(a)phénanthrén-17-ol, le 1-[2-(9 ou 10)-méthyl-9, 10-dihydrophénanthryl]-1-éthanol, le 2-hydroxy-9,10-dihydrophénanthrène, le 4'-oxo-7,8-cyclohexéno-9,10-dihydrophénanthrène-1-ol, le 4'-oxo-7,8-cyclohexénophénanthrén-1-ol, le 2-(2'-hydroxyéthyl)-9,10-dihydrophénanthrène et le 2-(2',3'-dihydroxyéthyl)-9,10-dihydrophénanthrène.
6. Procédé selon la revendication 1, qui comprend un élément choisi dans le groupe formé par :
le propionate de 4'-oxo-7,8-cyclohexano-9,10-dihydrophénanthrén-1-yle et l'acétate de 4'-oxo-7,8-cyclohexano-9,10-dihydrophénanthrén-1-yle.
7. Utilisation d'un antiandrogène comprenant un composé choisi dans le groupe formé par les composés de formule générale :



dans laquelle ledit cycle A est substitué ou non substitué et dans laquelle lorsque ledit cycle A est substitué, il est condensé en position 1-2 ou 2-3 avec un composé hydroxycycloalkyle ou cétycycloalkyle, ou il est substitué par un groupe hydroxyle, alkyle ou hydroxyalkyle en position 1, 2 ou 3,

dans laquelle ledit cycle B est saturé ou mono-insaturé en position 9-10 et est éventuellement substitué par un groupe alkyle inférieur en position 9 ou en position 10,

dans laquelle ledit cycle C est substitué ou non substitué, et dans laquelle lorsque ledit cycle C est substitué il est condensé en position 6-7 ou 7-8 avec un composé cétycycloalkyle ou hydroxycycloalkyle ; et

à condition que les cycles A et C ne puissent pas être l'un et l'autre non substitués, pour la préparation d'un médicament pour le traitement des troubles associés à une activité androgène excessive dans un organisme.

8. Utilisation selon la revendication 7, dans laquelle ledit antiandrogène est choisi dans le groupe formé par ;

le 1-[2-(9,10-dihydrophénanthryl)]-1-éthanol, le 6,7,16,17-tétrahydro-15H-cyclopenta[a]phénanthrén-17-ol, le 1-[2-(9 ou 10)-méthyl-9, 10-dihydrophénanthryl)]-1-éthanol, le 2-hydroxy-9,10-dihydrophénanthrène, la 6,7,16,17-tétrahydro-15H-cyclopenta[a]phénanthrén-17-one, le 4'-oxo-7,8-cyclohexéno-9,10-dihydrophénanthrén-1-ol, le 4'-oxo-7,8-cyclohexénophénanthrén-1-ol, le 2-(2'-hydroxyéthyl)-9,10-dihydrophénanthrène et le 2-(2', 3'-dihydroxyéthyl)-9,10-dihydrophénanthrène.




FIGURE 1




FIGURE 2




FIGURE 3




FIGURE 4